

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

I am informed from the office of the director that through mistake in cable dispatches there has been an error in recent reports from Piura, in that cases in Catacaos were accredited also to Piura, and therefore appeared twice in the statistics.

The 2 new cases here reported for Piura came from the neighboring village of Castilla.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

AFRICA—*Kimberly.*—Month ended January 11, 1908. Estimated population, 54,141. Total number of deaths, 119, including enteric fever 4, and 1 from scarlet fever.

AUSTRALIA—*New South Wales—Newcastle.*—Month of December, 1907. Estimated population, 51,423. Total number of deaths, 59, including enteric fever 2, and 1 from tuberculosis.

*Queensland—Brisbane.*—Month of December, 1907. Estimated population, 132,468. Total number of deaths, 153, including diphtheria 1, enteric fever 4, measles 1, and 8 from tuberculosis.

*State of Victoria, including the city of Melbourne.*—Quarter ended September 30, 1907. Estimated population, 1,237,998. Total number of deaths, 4,011.

AZORES—*St. Michaels and St. Marie.*—Month of December, 1907. Estimated population, 127,566. Total number of deaths, 252, including measles 9, enteric fever 9, scarlet fever 8, and 5 from tuberculosis.

CUBA.—Month of January, 1908. Reports from the 6 provinces, having an aggregate population of 1,572,797, show as follows: Deaths from contagious diseases: Diphtheria 11, leprosy 3, enteric fever 9, measles 3, yellow fever 6, and 115 from tuberculosis.

*Matanzas.*—Eleven days ended January 31, 1908. Estimated population, 35,515. Total number of deaths, 27, including whooping cough 1, and 5 from tuberculosis.

FRANCE—*Roubaix.*—Month of January, 1908. Estimated population, 121,115. Total number of deaths, 210, including 34 from tuberculosis.

*St. Etienne.*—Two weeks ended January 31, 1908. Estimated population, 150,000. Total number of deaths, 183, including enteric fever 1, diphtheria 5, measles 5, scarlet fever 2, and 24 from tuberculosis.

GREAT BRITAIN—*England and Wales.*—The deaths registered in 76 great towns in England and Wales during the week ended February 1, 1908, correspond to an annual rate of 18.2 per 1,000 population, which is estimated at 16,234,952.

*Dartmouth.*—Month of January, 1908. Estimated population, 7,000. Total number of deaths, 10. No deaths from contagious diseases reported.

*London.*—One thousand five hundred and eighty-four deaths were registered during the week, including measles 24, scarlet fever 12, diphtheria 24, enteric fever 1, whooping cough 43, tuberculosis 173, and 17 from diarrhea. The deaths from all causes correspond to an annual rate of 17.2 per 1,000. In Greater London 2,232 deaths were registered. In the "outer ring" the deaths included 5 from diphtheria, 3 from measles, 1 from scarlet fever, and 10 from whooping cough.

*Salford.*—Two weeks ended February 1, 1908. Estimated population, 236,670. Total number of deaths, 193, including diphtheria 2, measles 10, scarlet fever 2, whooping cough 3, and 12 from phthisis pulmonalis.

*Ireland.*—The average annual death rate represented by the deaths registered during the week ended February 1, 1908, in the 21 principal town districts of Ireland was 23.0 per 1,000 of the population, which is estimated at 1,131,959. The lowest rate was recorded in Drogheda, viz, 4.1 and the highest in Portadown, viz, 36.2 per 1,000.

*Scotland.*—The deaths registered in 8 principal towns during the week ended February 1, 1908, correspond to an annual rate of 22.3 per 1,000 of the population, which is estimated at 1,839,038. The highest rate of mortality was recorded in Greenock, viz, 25.4 and the lowest in Dundee, viz, 17.6 per 1,000. The aggregate number of deaths registered from all causes was 785, including diphtheria 6, enteric fever 2, measles 64, scarlet fever 7, and 26 from whooping cough.

JAMAICA—*Kingston.*—Month of January, 1908. Estimated population, 52,065. Total number of deaths, 123, including enteric fever 3, scarlet fever 1, and 16 from phthisis pulmonalis.

SPAIN—*Barcelona.*—Ten days ended January 20, 1908. Estimated population, 600,000. Total number of deaths, 431, including diphtheria 9, enteric fever 11, smallpox 1, and 32 from tuberculosis.

*Madrid.*—Month of January, 1908. Population, 556,663. Total number of deaths, 1,160, including diphtheria 2, enteric fever 11, measles 5, scarlet fever 3, and 125 from tuberculosis.

*Malaga.*—Month of November, 1907. Estimated population, 135,000. Total number of deaths, 325, including enteric fever 4, diphtheria 2, smallpox 15, and 33 from tuberculosis.

TURKEY—*Constantinople.*—Two weeks ended January 26, 1908. Estimated population 1,000,000. Total number of deaths, 545, including Asiatic cholera 8, diphtheria 5, enteric fever 14, measles 6, scarlet fever 6, smallpox 6, and 69 from tuberculosis.

URUGUAY—*Montevideo.*—Month of November, 1907. Estimated population, 309,511. Total number of deaths, 407, including diphtheria 5, enteric fever 3, scarlet fever 2, and 56 from tuberculosis.

WEST INDIES—*Curaçao.*—Two weeks ended February 7, 1908. Estimated population, 31,600. Total number of deaths, 10. No deaths from contagious diseases reported.